

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin, pan (Cocktail PAN-CK), 1mg/mL, Clone: [Cocktail PAN-CK], Mouse, Monoclonal BOT-BNUM0999-50

Artikelname	Cytokeratin, pan (Cocktail PAN-CK), 1mg/mL, Clone: [Cocktail PAN-CK], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0999-50
Hersteller Artikelnummer	BNUM0999-50
Alternativnummer	BOT-BNUM0999-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC
Spezies Reaktivität	Bovine, Canine, Gallus, Human, Monkey, Mouse, Rabbit, Rat
Immunogen	Human epidermal keratin
Produktbeschreibung	Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67 kDa (CK1), 64 kDa (CK3), 59 kDa (C...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[Cocktail PAN-CK]
Molekulargewicht	40-67 kDa (Multiple)

UniProt	Q7Z794
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user