

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD50 (ICAM-3)(ICAM3/1019), 1mg/mL, Clone: [ICAM3/1019], Mouse, Monoclonal BOT-BNUM1019-50

Artikelname	CD50 (ICAM-3)(ICAM3/1019), 1mg/mL, Clone: [ICAM3/1019], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1019-50
Hersteller Artikelnummer	BNUM1019-50
Alternativnummer	BOT-BNUM1019-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human ICAM3 protein
Produktbeschreibung	Recognizes an N-glycosylated glycoprotein of 120 kDa with intra-chain disulfide bonds, identified as CD50 or ICAM-3. CD50 is the major ligand for LFA-1 (CD11a/CD18) and may have signaling role to increase adhesion. It is expressed on thymocytes and T...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[ICAM3/1019]
Molekulargewicht	110-160 kDa

UniProt	P32942
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user