

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

PAX7 (PAX7/1187), 1mg/mL, Clone: [PAX7/1187], Mouse, Monoclonal BOT-BNUM1187-50

| | |
|--------------------------|---|
| Artikelname | PAX7 (PAX7/1187), 1mg/mL, Clone: [PAX7/1187], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM1187-50 |
| Hersteller Artikelnummer | BNUM1187-50 |
| Alternativnummer | BOT-BNUM1187-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment (aa300-600) of human PAX7 protein |
| Produktbeschreibung | The Pax gene family of nuclear transcription factors is comprised of nine members that function during embryogenesis to regulate the temporal and position-dependent differentiation of cells. In addition, the family is involved in a variety of signal ... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [PAX7/1187] |
| Molekulargewicht | 57 kDa |
| UniProt | P23759 |

| | |
|------------------------|---|
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris buffer with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |