

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

SUMO-1(SUMO1/1188), 1mg/mL, Clone: [SUMO1/1188], Mouse, Monoclonal BOT-BNUM1188-50

| | |
|--------------------------|---|
| Artikelname | SUMO-1(SUMO1/1188), 1mg/mL, Clone: [SUMO1/1188], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM1188-50 |
| Hersteller Artikelnummer | BNUM1188-50 |
| Alternativnummer | BOT-BNUM1188-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Rat |
| Immunogen | Recombinant human SUMO1 protein |
| Produktbeschreibung | This MAb is specific to SUMO-1 and shows no cross-reaction with either SUMO-2 or SUMO-3. The small ubiquitin-related modifier (SUMO) proteins, which include SUMO-1, SUMO-2 and SUMO-3, belong to the ubiquitin-like protein family. Like ubiquitin, the S... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [SUMO1/1188] |
| Molekulargewicht | 11.5 kDa (Monomer), 90 kDa (Heteromer) |

| | |
|------------------------|--|
| UniProt | P63165 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Predicted to show broad species reactivity Optimal dilution for a specific application should be determined by user |