

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

UGT1A9 (UDP glucuronosyltransferase 1 family, polypeptide A9)(UGT1A9/1229), 1mg/mL, Clone: [UGT1A9/1229], Mouse, Monoclonal BOT-BNUM1229-50

| | |
|--------------------------|---|
| Artikelname | UGT1A9 (UDP glucuronosyltransferase 1 family, polypeptide A9)(UGT1A9/1229), 1mg/mL, Clone: [UGT1A9/1229], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM1229-50 |
| Hersteller Artikelnummer | BNUM1229-50 |
| Alternativnummer | BOT-BNUM1229-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | A synthetic peptide specific to human UGT1A9 protein |
| Produktbeschreibung | This antibody recognizes a protein of about 60 kDa, which is identified as human UGT1A9. It does not cross-react with the other UGT1A isoforms including UGT1A7, UGT1A8, and UGT1A10 and shows a high degree of specificity. UGT1A9 is a UDP-glucuronosylt... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [UGT1A9/1229] |
| Molekulargewicht | 60 kDa |

| | |
|------------------------|--|
| UniProt | O60656 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |