

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Thymidine Phosphorylase / PD-ECGF (Angiogenesis Marker) (P-GF.44C), 1mg/mL, IgG1, Clone: [P-GF.44C], Mouse, Monoclonal BOT-BNUM1244-50

| | |
|--------------------------|---|
| Artikelname | Thymidine Phosphorylase / PD-ECGF (Angiogenesis Marker) (P-GF.44C), 1mg/mL, IgG1, Clone: [P-GF.44C], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM1244-50 |
| Hersteller Artikelnummer | BNUM1244-50 |
| Alternativnummer | BOT-BNUM1244-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human, Mouse, Rat |
| Immunogen | Recombinant full-length human Thymidine Phosphorylase (TP / PD-ECGF) protein |
| Produktbeschreibung | This antibody recognizes a protein (amino acid 482) of 55 kDa (in vivo 110 kDa homodimer), identified as platelet-derived endothelial growth factor (PD-ECGF), same as thymidine phosphorylase (TP) or gliostatin. In the presence of inorganic orthophosp... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [P-GF.44C] |

| | |
|------------------------|--|
| Molekulargewicht | 55 kDa |
| Isotyp | IgG1 |
| UniProt | P19971 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin) 1-2 ug/mL Western blotting 0.5-1 ug/mL Immunoprecipitation 0.5-1 ug/500ug protein lysate Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Optimal dilution for a specific application should be determined by user |