

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p57Kip2 (Mitotic Inhibitor/Suppressor Protein)(57P06), 1mg/mL, Clone: [57P06], Mouse, Monoclonal BOT-BNUM1264-50

Artikelname	p57Kip2 (Mitotic Inhibitor/Suppressor Protein)(57P06), 1mg/mL, Clone: [57P06], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1264-50
Hersteller Artikelnummer	BNUM1264-50
Alternativnummer	BOT-BNUM1264-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Mouse
Immunogen	Recombinant full-length human p57Kip2 protein
Produktbeschreibung	Recognizes a protein of 57 kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as a...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[57P06]
Molekulargewicht	57 kDa

UniProt	P49918
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 0.25-0.5 ug/mL for 30 minutes at RT Flow cytometry: 0.5-1 ug/million cells Immunofluorescence: 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user