

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Villin (GI-Mucosal & Urogenital Brush Border Marker) (VIL1/1314), 1mg/mL, Clone: [VIL1/1314], Mouse, Monoclonal BOT-BNUM1314-50

Artikelname	Villin (GI-Mucosal & Urogenital Brush Border Marker) (VIL1/1314), 1mg/mL, Clone: [VIL1/1314], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1314-50
Hersteller Artikelnummer	BNUM1314-50
Alternativnummer	BOT-BNUM1314-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human Villin fragment of 133 amino acid residues (aa179-311) (exact sequence is proprietary)
Produktbeschreibung	This antibody recognizes a protein of 95 kDa, which is identified as villin. It is a major constituent in the microvilli, which compose the brush border of epithelial cells forming absorptive surfaces of the intestinal and renal proximal tubular epit...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[VIL1/1314]

Molekulargewicht	93 kDa
UniProt	P09327
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Flow cytometry: 0.5-1 ug/million cells in 0.1mL Immunohistology (formalin): 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Western blotting 1-2 ug/mL Optimal dilution for a specific application should be determined by user