

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Protocadherin FAT2 (FAT Atypical Cadherin 2) (8C5), 1mg/mL, Clone: [8C5], Mouse, Monoclonal BOT-BNUM1320-50**

Artikelname	Protocadherin FAT2 (FAT Atypical Cadherin 2) (8C5), 1mg/mL, Clone: [8C5], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1320-50
Hersteller Artikelnummer	BNUM1320-50
Alternativnummer	BOT-BNUM1320-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Purified recombinant human FAT2 fusion protein
Produktbeschreibung	This antibody recognizes a protein of 480 kDa, which is identified as FAT2. The cadherins represent a family of Ca <sup>2+</sup> -dependent adhesion molecules that function to mediate cell-to-cell binding that is critical for the maintenance of structure and morp...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[8C5]
Molekulargewicht	480 kDa

UniProt	<a href="#">Q9NYQ8</a>
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user