

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CA19-9/Sialyl Lewisa (GI Tumor Marker) (CA19.9/1390R), 1mg/mL, Clone: [CA19.9/1390R], Rabbit, Monoclonal BOT-BNUM1390-50**

Artikelname	CA19-9/Sialyl Lewisa (GI Tumor Marker) (CA19.9/1390R), 1mg/mL, Clone: [CA19.9/1390R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM1390-50
Hersteller Artikelnummer	BNUM1390-50
Alternativnummer	BOT-BNUM1390-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Purified human CA19-9 protein
Produktbeschreibung	CA19-9, a carbohydrate epitope expressed on a high MW (>400 kDa) mucin glycoprotein, is a sialyl Lewisa structure which is synthesized from type 1 blood group precursor chains and is present in individuals expressing the Lewisa and/or Lewisb blood gr...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[CA19.9/1390R]
Molekulargewicht	>400 kDa

UniProt	Not Known
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence 0.5-1 ug/mL Immunohistology (formalin) 5-10 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user