

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

N-Cadherin / Cadherin-2 / CD325 (NCAD) (8C11), 1mg/mL, Clone: [8C11], Mouse, Monoclonal BOT-BNUM1426-50

Artikelname	N-Cadherin / Cadherin-2 / CD325 (NCAD) (8C11), 1mg/mL, Clone: [8C11], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1426-50
Hersteller Artikelnummer	BNUM1426-50
Alternativnummer	BOT-BNUM1426-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Mouse
Immunogen	Recombinant human N-cadherin extracellular domain (exact sequence is proprietary)
Produktbeschreibung	This antibody recognizes a protein of ~140 kDa, identified as N-Cadherin (NCAD), also known as CD325. NCAD is a member of the Cadherin superfamily, and consists of five extracellular repeats, a transmembrane domain and a cytoplasmic domain. CD325 def...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[8C11]

Molekulargewicht	130-140 kDa
UniProt	P19022
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user