

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Podoplanin (PDPN) (Lymphatic Endothelial & Mesothelial Marker) (PDPN/1433), 1mg/mL, Clone: [PDPN/1433], Mouse, Monoclonal BOT-BNUM1433-50

Artikelname	Podoplanin (PDPN) (Lymphatic Endothelial & Mesothelial Marker) (PDPN/1433), 1mg/mL, Clone: [PDPN/1433], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1433-50
Hersteller Artikelnummer	BNUM1433-50
Alternativnummer	BOT-BNUM1433-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human Podoplanin (PDPN) protein fragment (aa24-126) (exact sequence is proprietary)
Produktbeschreibung	This antibody recognizes a muco-protein of 38-43 kDa, which is identified Podoplanin (PDPN). It localizes in stromal cells of peripheral lymphoid tissue and thymic epithelial cells. As a regulator of the lymphatic endothelium, podoplanin probably pla...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[PDPN/1433]

Molekulargewicht	38-43 kDa
UniProt	Q86YL7
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Immunofluorescence 0.5-1 ug/mL Western blotting 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user