

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Desmoglein-3 (Squamous Cell Marker) (5G11), 1mg/mL, Clone: [5G11], Mouse, Monoclonal BOT-BNUM1439-50

Artikelname	Desmoglein-3 (Squamous Cell Marker) (5G11), 1mg/mL, Clone: [5G11], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1439-50
Hersteller Artikelnummer	BNUM1439-50
Alternativnummer	BOT-BNUM1439-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human DSG3 extracellular protein fragment (exact sequence is proprietary)
Produktbeschreibung	This antibody recognizes a protein of 130 kDa, identified as Desmoglein-3 (DSG3). This MAb is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein c...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[5G11]

Molekulargewicht	130 kDa
UniProt	P32926
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user