

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Alpha-1-Antichymotrypsin (SERPINA3) (Histiocytoma Marker) (AACT/1452), 1mg/mL, IgG1, Clone: [AACT/1452], Mouse, Monoclonal BOT-BNUM1452-50**

Artikelname	Alpha-1-Antichymotrypsin (SERPINA3) (Histiocytoma Marker) (AACT/1452), 1mg/mL, IgG1, Clone: [AACT/1452], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1452-50
Hersteller Artikelnummer	BNUM1452-50
Alternativnummer	BOT-BNUM1452-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human Antichymotrypsin (AACT) protein fragment (aa49-187) (exact sequence is proprietary)
Produktbeschreibung	Alpha-1 Antichymotrypsin (AACT) is a plasma protease inhibitor synthesized in the liver as a single glycopeptide chain. In human, the normal serum level of AACT is about one-tenth that of alpha-1-antitrypsin (AAT), with which it shares nucleic acid a...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[AACT/1452]

Molekulargewicht	65-76 kDa
Isotyp	IgG1
UniProt	<a href="#">P01011</a>
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Immunofluorescence 0.5-1 ug/mL Western blotting 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p>