

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

S100A4 (Marker of Tumor Metastasis) (S100A4/1482), 1mg/mL, Clone: [S100A4/1482], Mouse, Monoclonal BOT-BNUM1482-50

Artikelname	S100A4 (Marker of Tumor Metastasis) (S100A4/1482), 1mg/mL, Clone: [S100A4/1482], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1482-50
Hersteller Artikelnummer	BNUM1482-50
Alternativnummer	BOT-BNUM1482-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human, Mouse
Immunogen	Recombinant human S100A4 protein
Produktbeschreibung	S100A4 belongs to the S100 super-family of proteins containing 2 EF-hand calcium-binding domains. S100 genes include at least 25 members, including S100A1-S100A18, trichohyalin, filaggrin, repetin, S100P, and S100Z. S100A4 exerts its function via dir...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[S100A4/1482]
Molekulargewicht	10-12 kDa

UniProt	P26447
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user