

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Adipophilin / Perilipin-2 (Marker of Obesity) (ADFP/1493), 1mg/mL, Clone: [ADFP/1493], Mouse, Monoclonal BOT-BNUM1493-50

| | |
|--------------------------|---|
| Artikelname | Adipophilin / Perilipin-2 (Marker of Obesity) (ADFP/1493), 1mg/mL, Clone: [ADFP/1493], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM1493-50 |
| Hersteller Artikelnummer | BNUM1493-50 |
| Alternativnummer | BOT-BNUM1493-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human Adipophilin protein fragment (aa249-376) (exact sequence is proprietary) |
| Produktbeschreibung | Perilipin-2, also known as Adipophilin, belongs to the perilipin family, members of which coat intracellular lipid storage droplets. This 48 kDa protein is associated with the lipid globule surface membrane material, and maybe involved in development... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [ADFP/1493] |

| | |
|------------------------|---|
| Molekulargewicht | 48 kDa |
| UniProt | Q99541 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min ELISA For coating, order Ab without BSA Optimal dilution for a specific application should be determined by user |