

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD99 / MIC2 (Ewing s Sarcoma Marker) (MIC2/1495R), 1mg/mL, Clone: [MIC2/1495R], Rabbit, Monoclonal BOT-BNUM1495-50

Artikelname	CD99 / MIC2 (Ewing s Sarcoma Marker) (MIC2/1495R), 1mg/mL, Clone: [MIC2/1495R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM1495-50
Hersteller Artikelnummer	BNUM1495-50
Alternativnummer	BOT-BNUM1495-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human full-length MIC2 protein
Produktbeschreibung	This antibody recognizes a sialoglycoprotein of 27-32 kDa, identified as CD99, or MIC2 gene product, or E2 antigen. MIC2 gene is located in the pseudo-autosomal region of the human X and Y chromosome. MIC2 gene encodes two distinct proteins, which ar...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[MIC2/1495R]
Molekulargewicht	27-32 kDa

UniProt	P14209
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 2-4 ug/mL Flow cytometry 2-4 ug/million cells Immunohistology (formalin) 5-10 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Optimal dilution for a specific application should be determined by user