

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD56 / NCAM1 / NKH1 (Neuronal Cell Marker) (NCAM1/1496), 1mg/mL, Clone: [NCAM1/1496], Mouse, Monoclonal BOT-BNUM1496-50**

Artikelname	CD56 / NCAM1 / NKH1 (Neuronal Cell Marker) (NCAM1/1496), 1mg/mL, Clone: [NCAM1/1496], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1496-50
Hersteller Artikelnummer	BNUM1496-50
Alternativnummer	BOT-BNUM1496-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Spezies Reaktivität	Human
Immunogen	Recombinant human NCAM1 protein
Produktbeschreibung	This MAb reacts with an extracellular domain (close to transmembrane) of CD56/NCAM. Three isoforms of neural cell adhesion molecule (NCAM) are produced by differential splicing of the RNA transcript from a single gene. The 135 kDa isoform is the basi...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[NCAM1/1496]
Molekulargewicht	125 kDa 145 180

UniProt	<a href="#">P13591</a>
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry