

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD45 / LCA (B-Cell Marker) (F10-89-4), 1mg/mL, Clone: [F10-89-4], Mouse, Monoclonal BOT-BNUM1522-50**

|                          |   |
|--------------------------|---|
| Artikelname              | CD45 / LCA (B-Cell Marker) (F10-89-4), 1mg/mL, Clone: [F10-89-4], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNUM1522-50   |
| Hersteller Artikelnummer | BNUM1522-50   |
| Alternativnummer         | BOT-BNUM1522-50-50UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IF, IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Human T-Lymphocytes   |
| Produktbeschreibung      | CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms ar... |
| Klonalität               | Monoclonal  |
| Konzentration            | 1 mg/mL   |
| Klon-Bezeichnung         | [F10-89-4]  |

|                        |   |
|------------------------|---|
| Molekulargewicht       | 180-220 kDa   |
| UniProt                | <a href="#">P08575</a>  |
| Puffer                 | PBS, no BSA, no azide   |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |