

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Desmoglein-3 (Squamous Cell Marker) (DSG3/1535), 1mg/mL, Clone: [DSG3/1535], Mouse, Monoclonal BOT-BNUM1535-50

| | |
|--------------------------|---|
| Artikelname | Desmoglein-3 (Squamous Cell Marker) (DSG3/1535), 1mg/mL, Clone: [DSG3/1535], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM1535-50 |
| Hersteller Artikelnummer | BNUM1535-50 |
| Alternativnummer | BOT-BNUM1535-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human DSG3 extracellular protein fragment (exact sequence is proprietary) |
| Produktbeschreibung | This antibody recognizes a protein of 130 kDa, identified as Desmoglein-3 (DSG3). This MAb is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein c... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [DSG3/1535] |

| | |
|------------------------|---|
| Molekulargewicht | 130 kDa |
| UniProt | P32926 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |