

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Ep-CAM / CD326 (Epithelial Marker) (EGP40/1555R), 1mg/mL, Clone: [EGP40/1555R], Rabbit, Monoclonal BOT-BNUM1555-50

| | |
|--------------------------|---|
| Artikelname | Ep-CAM / CD326 (Epithelial Marker) (EGP40/1555R), 1mg/mL, Clone: [EGP40/1555R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNUM1555-50 |
| Hersteller Artikelnummer | BNUM1555-50 |
| Alternativnummer | BOT-BNUM1555-50-50UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | FC, IHC |
| Spezies Reaktivität | Human |
| Immunogen | A synthetic peptide (around aa 20-60) from the N-terminus of human TACSTD1 protein (exact sequence is proprietary) |
| Produktbeschreibung | This antibody recognizes a 40-43 kDa transmembrane epithelial glycoprotein, identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). Ep-CAM is expressed on baso-lateral cell surface in most simple epithelia ... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [EGP40/1555R] |

| | |
|------------------------|---|
| Molekulargewicht | 40-43 kDa |
| UniProt | P16422 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |