

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD162 (Selectin P Ligand) (PSGL1/1601), 1mg/mL, IgG1, Clone: [PSGL1/1601], Mouse, Monoclonal BOT-BNUM1601-50**

|                          |   |
|--------------------------|---|
| Artikelname              | CD162 (Selectin P Ligand) (PSGL1/1601), 1mg/mL, IgG1, Clone: [PSGL1/1601], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNUM1601-50   |
| Hersteller Artikelnummer | BNUM1601-50   |
| Alternativnummer         | BOT-BNUM1601-50-50UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant human full-length SELPLG (CD162) protein  |
| Produktbeschreibung      | CD162 glycoprotein functions as a high affinity counter-receptor for the cell adhesion molecules P-, E- and L- selectin expressed on myeloid cells and stimulated T lymphocytes. As such, this protein plays a critical role in leukocyte trafficking duri... |
| Klonalität               | Monoclonal  |
| Konzentration            | 1 mg/mL   |
| Klon-Bezeichnung         | [PSGL1/1601]  |
| Molekulargewicht         | 43-120 kDa  |

|                        |   |
|------------------------|---|
| Isotyp                 | IgG1  |
| UniProt                | <a href="#">Q14242</a>  |
| Puffer                 | PBS, no BSA, no azide   |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |