

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD27 (Tumor Necrosis Factor Receptor Superfamily 7) (LPFS2/1611), 1mg/mL, IgG1, Clone: [LPFS2/1611], Mouse, Monoclonal BOT-BNUM1611-50**

|                          |   |
|--------------------------|---|
| Artikelname              | CD27 (Tumor Necrosis Factor Receptor Superfamily 7) (LPFS2/1611),<br>1mg/mL, IgG1, Clone: [LPFS2/1611], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNUM1611-50   |
| Hersteller Artikelnummer | BNUM1611-50   |
| Alternativnummer         | BOT-BNUM1611-50-50UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF, IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant human CD22 protein (exact sequence is proprietary)  |
| Produktbeschreibung      | This antibody recognizes a protein of a disulfide-linked 120 kDa dimer, identified as CD27. It is expressed on the majority of peripheral T cells, medullary thymocytes, memory-type B cells, and natural killer cells. It is a transmembrane phosphoglyco... |
| Klonalität               | Monoclonal  |
| Konzentration            | 1 mg/mL   |
| Klon-Bezeichnung         | [LPFS2/1611]  |
| Molekulargewicht         | 120 kDa   |

|                        |   |
|------------------------|---|
| Isotyp                 | IgG1  |
| UniProt                | <a href="#">P26842</a>  |
| Puffer                 | PBS, no BSA, no azide   |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris buffer with 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |