

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD79a (IGA/1688R), 1mg/mL, Clone: [IGA/1688R], Rabbit, Monoclonal BOT-BNUM1688-50

Artikelname	CD79a (IGA/1688R), 1mg/mL, Clone: [IGA/1688R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM1688-50
Hersteller Artikelnummer	BNUM1688-50
Alternativnummer	BOT-BNUM1688-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79a protein.
Produktbeschreibung	A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[IGA/1688R]
Molekulargewicht	44 kDa

UniProt	P11912
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user