

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

S100B (Astrocyte and Melanoma Marker) (S100B/1706R), 1mg/mL, Clone: [S100B/1706R], Rabbit, Monoclonal BOT-BNUM1706-50

Artikelname	S100B (Astrocyte and Melanoma Marker) (S100B/1706R), 1mg/mL, Clone: [S100B/1706R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM1706-50
Hersteller Artikelnummer	BNUM1706-50
Alternativnummer	BOT-BNUM1706-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Bovine, Human, Mouse, Rat
Immunogen	Recombinant full-length human S100B protein
Produktbeschreibung	S100A and S100B proteins are two members of the S100 family of calcium binding proteins. S100A is composed of an alpha and a beta chain whereas S100B is composed of two beta chains. This antibody is specific against an epitope located on the beta-cha...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[S100B/1706R]
Molekulargewicht	10-12 kDa

UniProt	P04271
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user