

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

AMACR / p504S (Prostate Cancer Marker) (AMACR/1723), 1mg/mL, Clone: [AMACR/1723], Mouse, Monoclonal BOT-BNUM1723-50

| | |
|--------------------------|---|
| Artikelname | AMACR / p504S (Prostate Cancer Marker) (AMACR/1723), 1mg/mL, Clone: [AMACR/1723], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM1723-50 |
| Hersteller Artikelnummer | BNUM1723-50 |
| Alternativnummer | BOT-BNUM1723-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human full-length AMACR protein |
| Produktbeschreibung | This antibody recognizes a protein of 42 kDa, which is identified as Alpha-methylacyl-CoA Racemase (AMACR), also known as p504S. It is an enzyme that is involved in bile acid biosynthesis and beta-oxidation of branched-chain fatty acids. AMACR is ess... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [AMACR/1723] |
| Molekulargewicht | 54 kDa |

| | |
|------------------------|---|
| UniProt | Q9UHK6 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Optimal dilution for a specific application should be determined by user |