

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Glycophorin A / CD235a (Erythrocyte Marker) (GYPA/1725R), 1mg/mL, Clone: [GYPA/1725R], Rabbit, Monoclonal BOT-BNUM1725-50**

Artikelname	Glycophorin A / CD235a (Erythrocyte Marker) (GYPA/1725R), 1mg/mL, Clone: [GYPA/1725R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM1725-50
Hersteller Artikelnummer	BNUM1725-50
Alternativnummer	BOT-BNUM1725-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human full-length Glycophorin A protein
Produktbeschreibung	This antibody recognizes a sialoglycoprotein of 39 kDa, identified as glycophorin A (GPA). It is present on red blood cells (RBC) and erythroid precursor cells. It has been shown that glycophorin acts as the receptor for Sandei virus and parvovirus. ...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[GYPA/1725R]
Molekulargewicht	39 kDa

UniProt	<a href="#">P02724</a>
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user