

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Chromogranin A / CHGA (Neuroendocrine Marker) (CHGA/1731R), 1mg/mL, Clone: [CHGA/1731R], Rabbit, Monoclonal BOT-BNUM1731-50

| | |
|--------------------------|---|
| Artikelname | Chromogranin A / CHGA (Neuroendocrine Marker) (CHGA/1731R), 1mg/mL, Clone: [CHGA/1731R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNUM1731-50 |
| Hersteller Artikelnummer | BNUM1731-50 |
| Alternativnummer | BOT-BNUM1731-50-50UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human full-length Chromogranin A protein |
| Produktbeschreibung | Chromogranin A is present in neuroendocrine cells throughout the body, including the neuroendocrine cells of the large and small intestine, adrenal medulla and pancreatic islets. It is an excellent marker for carcinoid tumors, pheochromocytomas, para... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [CHGA/1731R] |
| Molekulargewicht | 68-75 kDa |

| | |
|------------------------|--|
| UniProt | P10645 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |