

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Glycophorin A / CD235a (Erythrocyte Marker) (JC159), 1mg/mL, Clone: [JC159], Mouse, Monoclonal BOT-BNUM1737-50**

Artikelname	Glycophorin A / CD235a (Erythrocyte Marker) (JC159), 1mg/mL, Clone: [JC159], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1737-50
Hersteller Artikelnummer	BNUM1737-50
Alternativnummer	BOT-BNUM1737-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Formalin fixed membrane from a Hairy cell leukemia
Produktbeschreibung	This antibody recognizes a sialoglycoprotein of 39 kDa, identified as glycophorin A (GPA). It is present on red blood cells (RBC) and erythroid precursor cells. It has been shown that glycophorin acts as the receptor for Sandei virus and parvovirus. ...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[JC159]
Molekulargewicht	39 kDa

UniProt	<a href="#">P02724</a>
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user