

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **MyoD1 (Rhabdomyosarcoma Marker) (MYOD1/2075R), 1mg/mL, Clone: [MYOD1/2075R], Rabbit, Monoclonal BOT-BNUM2075-50**

Artikelname	MyoD1 (Rhabdomyosarcoma Marker) (MYOD1/2075R), 1mg/mL, Clone: [MYOD1/2075R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM2075-50
Hersteller Artikelnummer	BNUM2075-50
Alternativnummer	BOT-BNUM2075-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human MyoD1 protein
Produktbeschreibung	This antibody recognizes a phosphoprotein of 45 kDa, identified as MyoD1. This MAb does not cross react with myogenin, Myf5, or Myf6. Antibody to MyoD1 labels the nuclei of myoblasts in developing muscle tissues. MyoD1 is not detected in normal adult...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[MYOD1/2075R]
Molekulargewicht	45 kDa

UniProt	<a href="#">P15172</a>
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user