

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Ubiquitin (Autophagy Marker) (UBB/2122), 1mg/mL, Clone: [UBB/2122], Mouse, Monoclonal BOT-BNUM2122-50

Artikelname	Ubiquitin (Autophagy Marker) (UBB/2122), 1mg/mL, Clone: [UBB/2122], Mouse, Monoclonal
Artikelnummer	BOT-BNUM2122-50
Hersteller Artikelnummer	BNUM2122-50
Alternativnummer	BOT-BNUM2122-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant fragment of human Ubiquitin protein (around aa 1-119) (exact sequence is proprietary)
Produktbeschreibung	Ubiquitin is a highly conserved and plays an essential role in the ubiquitin-proteasome pathway. In ubiquitination process, it is first activated by forming a thiol-ester complex with the activation component E1, which is then transferred to ubiquiti...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[UBB/2122]

Molekulargewicht	9 kDa
UniProt	<a href="#">P0CG47</a>
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes ELISA For coating, order Ab without BSA Optimal dilution for a specific application should be determined by user