

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mesothelin (Mesothelial Marker) (MSLN/2131), 1mg/mL, Clone: [MSLN/2131], Mouse, Monoclonal BOT-BNUM2131-50

Artikelname	Mesothelin (Mesothelial Marker) (MSLN/2131), 1mg/mL, Clone: [MSLN/2131], Mouse, Monoclonal
Artikelnummer	BOT-BNUM2131-50
Hersteller Artikelnummer	BNUM2131-50
Alternativnummer	BOT-BNUM2131-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Recombinant fragment (around aa 273-407) of human Mesothelin (MSLN) protein (exact sequence is proprietary)
Produktbeschreibung	Mesothelin is a 40 kDa glycosyl-phosphatidylinositol-anchored glycoprotein cleaved from a 69 kDa precursor protein. Mesothelin immunoreactivity is high in cancers of the ovary (serous papillary, endometrioid and undifferentiated) and pancreas, with I...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[MSLN/2131]

Molekulargewicht	40 kDa (Mature form), 69 kDa (Precursor)
UniProt	Q13421
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user