

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TACSTD2 / TROP2 (Epithelial Marker) (TACSTD2/2153), 1mg/mL, Clone: [TACSTD2/2153], Mouse, Monoclonal BOT-BNUM2153-50

Artikelname	TACSTD2 / TROP2 (Epithelial Marker) (TACSTD2/2153), 1mg/mL, Clone: [TACSTD2/2153], Mouse, Monoclonal
Artikelnummer	BOT-BNUM2153-50
Hersteller Artikelnummer	BNUM2153-50
Alternativnummer	BOT-BNUM2153-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant fragment of human TACSTD2 protein (around aa 31-274) (exact sequence is proprietary)
Produktbeschreibung	TACSTD2 is a cell surface glycoprotein receptor. It is a single pass type I membrane protein containing one thyroglobulin type-1 domain, an epidermal growth factor-like repeat, a phosphatidylinositol binding site and tyrosine phosphorylation sites ne...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[TACSTD2/2153]

Molekulargewicht	40 kDa
UniProt	P09758
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user