

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **HSP60 (Heat Shock Protein 60) (Mitochondrial Marker) (rGROEL/780), 1mg/mL, Clone: [rGROEL/780], Mouse, Monoclonal BOT-BNUM2207-50**

|                          |   |
|--------------------------|---|
| Artikelname              | HSP60 (Heat Shock Protein 60) (Mitochondrial Marker) (rGROEL/780), 1mg/mL, Clone: [rGROEL/780], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNUM2207-50   |
| Hersteller Artikelnummer | BNUM2207-50   |
| Alternativnummer         | BOT-BNUM2207-50-50UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC, WB   |
| Spezies Reaktivität      | Bovine, Canine, Gallus, Hamster, Human, Invertebrate, Monkey, Mouse, Porcine, Rabbit, Rat, Sheep, Xenopus   |
| Immunogen                | Recombinant full-length human HSP60 protein   |
| Produktbeschreibung      | This antibody recognizes a 60 kDa protein, identified as the heat shock protein 60 (hsp60). Its epitope is localized between aa 383-447 of human hsp60. A wide variety of environmental and pathophysiological stressful conditions trigger the synthesis ... |
| Klonalität               | Monoclonal  |
| Konzentration            | 1 mg/mL   |
| Klon-Bezeichnung         | [rGROEL/780]  |

|                        |   |
|------------------------|---|
| Molekulargewicht       | 60 kDa  |
| UniProt                | <a href="#">P10809</a>  |
| Puffer                 | PBS, no BSA, no azide   |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.25-0.5 ug/mL Optimal dilution for a specific application should be determined by user |