

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **NKX2.2 (Neuroendocrine & Ewing s Sarcoma Marker) (rNX2/1523), 1mg/mL, IgG1, Clone: [rNX2/1523], Mouse, Monoclonal BOT-BNUM2309-50**

Artikelname	NKX2.2 (Neuroendocrine & Ewing s Sarcoma Marker) (rNX2/1523), 1mg/mL, IgG1, Clone: [rNX2/1523], Mouse, Monoclonal
Artikelnummer	BOT-BNUM2309-50
Hersteller Artikelnummer	BNUM2309-50
Alternativnummer	BOT-BNUM2309-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Human recombinant NKX2.2 protein fragment (around aa1-119) (exact sequence is proprietary)
Produktbeschreibung	Expression of NKX2.2 has been found in neuroendocrine tumors of the gut, making it a potential marker for the study of gastrointestinal neuroendocrine tumors. More recently, NKX2.2 protein was identified as a target of EWS-FLI-1, the fusion protein s...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[rNX2/1523]

Molekulargewicht	40-50 kDa
Isotyp	IgG1
UniProt	<a href="#">O95096</a>
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user