

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

SDHB (Succinate Dehydrogenase B) (Pheochromocytoma Marker)(SDHB/2382), 1mg/mL, Clone: [SDHB/2382], Mouse, Monoclonal BOT-BNUM2382-50

| | |
|--------------------------|---|
| Artikelname | SDHB (Succinate Dehydrogenase B) (Pheochromocytoma Marker)(SDHB/2382), 1mg/mL, Clone: [SDHB/2382], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUM2382-50 |
| Hersteller Artikelnummer | BNUM2382-50 |
| Alternativnummer | BOT-BNUM2382-50-50UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment (around aa 165-273) of human SDHB protein (exact sequence is proprietary) |
| Produktbeschreibung | Succinate dehydrogenase (SDH) is Complex II in the mitochondria, vital for mitochondrial electron transport, as well as Krebs cycle function. Four subunits comprise the SDH protein complex: a flavochrome subunit (SDHA), an iron-sulfur protein (SDHB) ... |
| Klonalität | Monoclonal |
| Konzentration | 1 mg/mL |
| Klon-Bezeichnung | [SDHB/2382] |
| Molekulargewicht | 32 kDa |

| | |
|------------------------|--|
| UniProt | P21912 |
| Puffer | PBS, no BSA, no azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |