

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD5 (Mantle Cell Lymphoma Marker) (CD5/2416), 1mg/mL, Clone: [CD5/2416], Mouse, Monoclonal BOT-BNUM2416-50**

Artikelname	CD5 (Mantle Cell Lymphoma Marker) (CD5/2416), 1mg/mL, Clone: [CD5/2416], Mouse, Monoclonal
Artikelnummer	BOT-BNUM2416-50
Hersteller Artikelnummer	BNUM2416-50
Alternativnummer	BOT-BNUM2416-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant fragment of human CD5 protein (around aa 269-366) (exact sequence is proprietary)
Produktbeschreibung	This antibody recognizes a 67 kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95% of thymocytes and 72% of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas. Anti-CD5 is a...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[CD5/2416]

Molekulargewicht	67 kDa
UniProt	<a href="#">P06127</a>
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Western Blot 0.5-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user