

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

BOB.1 (B-Cell Marker) (BOB1/2424), 1mg/mL, Clone: [BOB1/2424], Mouse, Monoclonal BOT-BNUM2424-50

Artikelname	BOB.1 (B-Cell Marker) (BOB1/2424), 1mg/mL, Clone: [BOB1/2424], Mouse, Monoclonal
Artikelnummer	BOT-BNUM2424-50
Hersteller Artikelnummer	BNUM2424-50
Alternativnummer	BOT-BNUM2424-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human BOB1 (POU2AF1) protein fragment (around aa 148-255) (exact sequence is proprietary)
Produktbeschreibung	BOB-1 is a transcriptional coactivator that specifically associates with either POU2F1/OCT1 or POU2F2/OCT2, boosting the promoter activity of those complexes. It is essential for the response of B-cells to antigens, required for the formation of germ...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[BOB1/2424]

Molekulargewicht	35 kDa
UniProt	Q16633
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user