

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p63 (Squamous, Basal & Myoepithelial Cell Marker) (TP63/2427), 1mg/mL, Clone: [TP63/2427], Mouse, Monoclonal BOT-BNUM2427-50

Artikelname	p63 (Squamous, Basal & Myoepithelial Cell Marker) (TP63/2427), 1mg/mL, Clone: [TP63/2427], Mouse, Monoclonal
Artikelnummer	BOT-BNUM2427-50
Hersteller Artikelnummer	BNUM2427-50
Alternativnummer	BOT-BNUM2427-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human p63 protein
Produktbeschreibung	p63 is a homolog of the tumor suppressor p53. It is identified in basal cells in the epithelial layers of a variety of tissues, including epidermis, cervix, urothelium, breast and prostate. p63 was detected in nuclei of the basal epithelium in normal...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[TP63/2427]
Molekulargewicht	63 kDa

UniProt	Q9H3D4
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user