

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Ki-67 (Proliferating Cell Marker) (MKI67/2462), 1mg/mL, Clone: [MKI67/2462], Mouse, Monoclonal BOT-BNUM2462-50**

Artikelname	Ki-67 (Proliferating Cell Marker) (MKI67/2462), 1mg/mL, Clone: [MKI67/2462], Mouse, Monoclonal
Artikelnummer	BOT-BNUM2462-50
Hersteller Artikelnummer	BNUM2462-50
Alternativnummer	BOT-BNUM2462-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human Ki67 protein fragment (around aa 2293-2478) (exact sequence is proprietary)
Produktbeschreibung	Ki-67 antigen is a nuclear, non-histone protein that is present in all stages of the cell cycle except G0. This characteristic makes Ki-67 an excellent marker for proliferating cells and is commonly used as one of the prognostic factors in cancer stu...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[MKI67/2462]

Molekulargewicht	345 kDa and 395 kDa
UniProt	<a href="#">P46013</a>
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user