

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Histone Deacetylase 7 (Phospho-S155) polyclonal antibody, Unconjugated, Rabbit BWT-BS64588**

|                          |   |
|--------------------------|---|
| Artikelname              | Histone Deacetylase 7 (Phospho-S155) polyclonal antibody, Unconjugated, Rabbit  |
| Artikelnummer            | BWT-BS64588   |
| Hersteller Artikelnummer | BS64588   |
| Alternativnummer         | BWT-BS64588-50UL,BWT-BS64588-100UL  |
| Hersteller               | Bioworld Technology   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | IF, WB  |
| Spezies Reaktivität      | Human, Mouse, Rat   |
| Immunogen                | Synthetic phosphopeptide derived from human Histone Deacetylase 7 around the phosphorylation site of Serine 155.  |
| Konjugation              | Unconjugated  |
| Produktbeschreibung      | In the intact cell, DNA closely associates with histones and other nuclear proteins to form chromatin. The remodeling of chromatin is believed to be a critical component of transcriptional regulation and a major source of this remodeling is brought a... |
| Molekulargewicht         | ~ 105 kDa   |
| UniProt                  | <a href="#">Q8WUI4</a>  |

|                        |   |
|------------------------|---|
| Reinheit               | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE). |
| Formulierung           | Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2  |
| Application Verdünnung | WB:1:500~1:1000 IF:1:50~1:200   |
| Anwendungsbeschreibung | Histone Deacetylase 7 (Phospho-S155) polyclonal antibody detects endogenous levels of Histone Deacetylase 7 protein only when phosphorylated at Ser155.     |