

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Recombinant Human Endothelial-MonocyteAActivating Polypeptide II (rHuEMAP-II) BWT-PR1015

Artikelname	Recombinant Human Endothelial-MonocyteAActivating Polypeptide II (rHuEMAP-II)
Artikelnummer	BWT-PR1015
Hersteller Artikelnummer	PR1015
Alternativnummer	BWT-PR1015-5UG,BWT-PR1015-20UG,BWT-PR1015-1.0MG
Hersteller	Bioworld Technology
Kategorie	Proteine/Peptide
Produktbeschreibung	EMAP-II is a tumor derived cytokine that exerts a wide range of activities on endothelial cells, monocytes and neutrophils. EMAP-II inhibits endothelial cell proliferation, vasculogenesis, neovessel formation, and can induce apoptosis. It is also che...
Molekulargewicht	Approximately 18.3 KDa, a single non-glycosylated polypeptide chain containing 166 amino acids.
Quelle	Escherichia coli.
Reinheit	>98% by SDS-PAGE and HPLC analyses.
Formulierung	Sterile Filtered White lyophilized (freeze-dried) powder.
Sequenz	SKPIDVSRDL LRIGCIITAR KHPDADSLYV EEVDVGEIAP RTVVSGLVNH VPLEQMQRNM VILLCNLKPA KMRGVLSQAM VMCASSPEKI EILAPPNGSV PGDRITFDAF PGEPDKELNP KKKIWEQIQP DLHTNDECVA TYKGVPFVVK GKGVCRAQTM SNSGIK

Formel	Lyophilized from a 0.2mm filtered concentrated solution in 20mM PB, pH 7.4, 130mM NaCl.
Anwendungsbeschreibung	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at <-20C. Further dilutions should be made in appropriate buffered solutions.