

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Anti-Plasminogen PLG Antibody Peroxidase Conjugated, HRP, Goat, Polyclonal BOB-H00702

Artikelname	Anti-Plasminogen PLG Antibody Peroxidase Conjugated, HRP, Goat, Polyclonal
Artikelnummer	BOB-H00702
Hersteller Artikelnummer	H00702
Alternativnummer	BOB-H00702-100UG
Hersteller	Boster Bio
Wirt	Goat
Kategorie	Antikörper
Applikation	ELISA, IHC, WB
Spezies Reaktivität	Human
Immunogen	Plasminogen [Human Plasma]
Konjugation	HRP
Produktbeschreibung	Boster Bio Anti-Plasminogen PLG Antibody Peroxidase Conjugated (Catalog H00702). Tested in ELISA, WB applications. This antibody reacts with Human....
Klonalität	Polyclonal
Konzentration	1.0 mg/mL by UV absorbance at 280 nm
Molekulargewicht	90569 MW

UniProt	P00747
Puffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free, 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Reinheit	Anti-Plasminogen is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum as well as purified and partially purified Plasminogen [Human Plasma]. Cross-reactivity against Plasminogen from other sources is unknown.
Formulierung	Lyophilized
Application Verdünnung	ELISA: 1:2,000 - 1:10,000IHC: 1:250 - 1:1,000WB: 1:500 - 1:2,500Anti-Plasminogen has been tested by ELISA and western blot. This product is assayed against 1.0 µg of Plasminogen [Human Plasma] in a standard capture ELISA using ABTS (2,2-azino-bis-[3-ethyl
Anwendungsbeschreibung	Anti-Plasminogen has been tested by ELISA and western blot. This product is assayed against 1.0 µg of Plasminogen [Human Plasma] in a standard capture ELISA using ABTS (2,2-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) as a substrate for 30 minutes