

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mucin-1 Antibody, Clone: [SPM533], Mouse, Monoclonal NSJ-V9048-20UG

Artikelname	Mucin-1 Antibody, Clone: [SPM533], Mouse, Monoclonal
Artikelnummer	NSJ-V9048-20UG
Hersteller Artikelnummer	V9048-20UG
Alternativnummer	NSJ-V9048-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Delipidated extract of human milk fat globule membranes was used as the immunogen for this Mucin-1 antibody.
Produktbeschreibung	Mucin-1 is a 265-400kDa glycoprotein. This protein is proteolytically cleaved into alpha and beta subunits that form a heterodimeric complex. The N-terminal alpha subunit and the C-terminal beta subunit. . The alpha subunit has cell adhesive properti...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM533]
UniProt	P15941
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	The optimal dilution of the Mucin-1 antibody for each application should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 minutes.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.