

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat Anti-Guinea Pig IgG F(ab)2 Antibody Biotin Conjugated - 606-1604, Polyclonal DNA-SEC-182916

Article Name	Goat Anti-Guinea Pig IgG F(ab)2 Antibody Biotin Conjugated - 606-1604, Polyclonal
Biozol Catalog Number	DNA-SEC-182916
Supplier Catalog Number	DNA-SEC-182916
Alternative Catalog Number	DNA-SEC-182916
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	ELISA,IHC,WB
Species Reactivity	Guinea pig
Immunogen	Guinea Pig IgG F(ab)2 fragment
Conjugation	Biotin
Format	IgG
Target Specificity	IgG (F(ab')2)
Cross-Adsorption (MinX)	no cross-adsorbtion
Product Description	Anti-Guinea Pig IgG F(ab)2 Biotin Antibody generated in goat is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlled conditions of temperature, time and pH. F(ab)2 molecules lack the F...

Clonality	Polyclonal
Concentration	2.0 mg/mL
Isotype	Ig
Buffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Guinea Pig IgG, Guinea Pig IgG F(ab)2 and Guinea Pig Serum. No reaction was observed against Guinea Pig IgG F(c).
Form	Lyophilized
Formula	10 mM NaPO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target	Guinea Pig
Antibody Type	Secondary Antibody
Application Dilute	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	This product has been assayed against 1.0 ug of Guinea Pig IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin S000-03 and ABTS (2,2-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:28,000 to 1:114,000 of the reconstitution concentration is suggested for this product.