

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Human BMP4 protein, His tag, Unconjugated GTX00079-PRO**

Artikelname	Human BMP4 protein, His tag, Unconjugated
Artikelnummer	GTX00079-PRO
Hersteller Artikelnummer	GTX00079-pro
Alternativnummer	GTX00079-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	<a href="#">652</a>
UniProt	<a href="#">P12644</a>
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Ser22~Arg408 (NP_001193.2)

#### Anwendungsbeschreibung

BMP4 (Bone morphogenetic protein 4) is a member of the bone morphogenetic protein family, which is involved in bone and cartilage development, specifically tooth and limb development and fracture repair. It has been proven that HJV (Hemojuvelin) acts as a coreceptor of BMPs, including BMP4, therefore, a binding ELISA assay was constructed to detect the association of recombinant human BMP4 with recombinant human HJV. Briefly, BMP4 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100  $\mu$ l were then transferred to HJV-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-BMP4 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50  $\mu$ l stop solution to the wells and read at 450nm immediately. The binding activity of BMP4 with HJV was in a dose dependent manner.