

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Human TGF beta 1 protein, His tag, Unconjugated GTX00101-PRO

Artikelname	Human TGF beta 1 protein, His tag, Unconjugated
Artikelnummer	GTX00101-PRO
Hersteller Artikelnummer	GTX00101-pro
Alternativnummer	GTX00101-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	<a href="#">7040</a>
UniProt	<a href="#">P01137</a>
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Ala279~Ser390 (NP_000651.3)

Anwendungsbeschreibung

TGF-beta1 (Transforming growth factor beta 1) is a multifunctional set of peptides that controls proliferation, differentiation, and other functions in many cell types. TGF beta 1 has been shown to interact with TGF beta receptor 1, Decorin, LTBP1 and so on. Thus we have conducted a binding ELISA assay to detect the interaction of recombinant human TGF-beta1 with recombinant human LTBP1. Briefly, TGF-beta1 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100  $\mu$ l were then transferred to LTBP1-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-LTBP1 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50  $\mu$ l stop solution to the wells and read at 450nm immediately. The binding activity of TGF-beta1 with LTBP1 was in a dose dependent manner.