

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## **Product Datasheet**

## Human CX3CL1 protein, His tag (active), Unconjugated GTX00214-PRO

Artikelname	Human CX3CL1 protein, His tag (active), Unconjugated
Artikelnummer	GTX00214-PRO
Hersteller Artikelnummer	GTX00214-pro
Alternativnummer	GTX00214-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	6376
UniProt	P78423
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Gln25~Pro332 (NP_002987.1)

is a large cytokine protein of 373 amino acids, it contains multiple domains and is the only known member of the CX3C chemokine family. Soluble CX3CL1 potently chemoattracts T cells and monocytes, while the cell-bound chemokine promotes strong adhesion of leukocytes to activated endothelial cells, where it is primarily expressed. Thus, chemotaxis assay used 24-well microchemotaxis system was undertaken to detect the chemotactic effect of CX3CL1 on the human monocytic cell line THP-1. Briefly, THP-1 cells were seeded into the upper chambers (150  $\mu$ l cell suspension, 1x106 cells/ml in RPMI-1640 with FBS free) and SLC (1 ng/ml, 10 ng/ml, 100 ng/ml and 1000 ng/ml diluted separately in serum free RPMI-1640 ) was added in lower chamber with a polycarbonate filter (8um pore size) used to separate the two compartments. After incubation at 37C with 5% CO2 for 1h, the filter was removed, then cells in low chamber were observed by inverted microscope at low magnification (\*100) and the number of migrated

cells were counted at high magnification (\*400) randomly. And

CX3CL1 is able to induce migration of THP-1 cells.

Chemokine C-X3-C-Motif Ligand 1 (CX3CL1) also known as fractalkine

Anwendungsbeschreibung