

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human CHI3L1 protein, His tag (active), Unconjugated GTX00219-PRO

Artikelname	Human CHI3L1 protein, His tag (active), Unconjugated
Artikelnummer	GTX00219-PRO
Hersteller Artikelnummer	GTX00219-pro
Alternativnummer	GTX00219-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	1116
UniProt	P36222
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Asn112~Thr237+Ile299~Lys377 (NP_001267.2)

Anwendungsbeschreibung

Cell proliferation of Raji cells after stimulated with GP39. Cell proliferation of Raji cells after stimulated with GP39. Glycoprotein 39, Cartilage (GP39) also known as Chitinase-3-like protein 1 (CHI3L1) is a secreted glycoprotein that is approximately 40kDa in size that in humans is encoded by the CHI3L1 gene. GP39 plays a role in cancer cell proliferation, survival, invasiveness and in the regulation of cellmatrix interactions. To test the effect of GP39 on cell proliferation, Raji cells were seeded into triplicate wells of 96-well plates at a density of 5000 cells/well with 1% serum standard RPMI-1640 which contains various concentrations of recombinant human GP39. After incubated for 5 days, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10 µl of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating the plate for 1-4 hours at 37C. Proliferation of Raji cells after incubation with GP39 for 5 days observed by inverted microscope.