

Diagnostica Vertrieb GmbH, Oehleckerring 11-13

22419 Hamburg, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human FGF6 protein, His and GST tag, Unconjugated GTX00224-PRO

Artikelname	Human FGF6 protein, His and GST tag, Unconjugated
Artikelnummer	GTX00224-PRO
Hersteller Artikelnummer	GTX00224-pro
Alternativnummer	GTX00224-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	2251
UniProt	P10767
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His and GST-Tag, Gly41~Ile208 (NP_066276.2)

Anwendungsbeschreibung

Fibroblast growth factor 6 (FGF6) is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. It may also play a importent role in muscle regeneration or differentiation. Besides, Fibroblast Growth Factor Receptor 1 (FGFR1) has been identified as an interactor of FGF6, thus a binding ELISA assay was conducted to detect the interaction of recombinant human FGF6 and recombinant human FGFR1. Briefly, FGF6 were diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100 μ l were then transferred to FGFR1-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-FGF6 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of FGF6 and FGFR1 was in a dose dependent manner.